

Copper 20g



Distributed by: **Pacific Biologics Pty. Ltd.**, 35 Beach St,  
Kippa Ring QLD 4021 Ph: (07) 3283 5077 www.pacfbio.com.au  
Manufactured by: Agrimin Ltd, UK.  
APVMA Approval No. 81699/103765  
Expiry Date:  
Batch No.:  
Cu20gAUSI - 16/7

Protect from light.  
Store below 25°C (air conditioning). Store in original container.

**STORAGE**  
Dispose of container by wrapping with paper and putting in garbage.

**DISPOSAL**  
This product is contraindicated for use in cattle suffering from jaundice and other liver disorders such as liver fluke, or stock grazing plants containing toxic alkaloids, e.g. Heliotrope, Ragwort, Patersons Curse (Salvation Jane).  
Seek the advice of a veterinary or cattle husbandry specialist to ascertain copper deficiency and type of treatment required for cattle under consideration.

**Dosage and Administration**  
Only repeat treatment within 12 months when specifically indicated by laboratory tests. Administer orally. Place the bolus at the back of the mouth using a suitable applicator. It is important that the applicator's instructions are strictly followed. One bolus should be administered per animal over 200 kg. One bolus per 200 kg.

**General Directions**  
To be effective, the bolus must be swallowed by the animal.  
Dose pregnant females early in pregnancy to allow protection of the calf against copper deficiency.

**Contraindications**  
SHOULD NOT BE ADMINISTERED TO CATTLE LESS THAN 5 MONTHS OF AGE OR 200 kg BODYWEIGHT  
Excessive copper is toxic. This product is contraindicated for use unless copper deficiency is confirmed.

**DIRECTIONS FOR USE**  
AGRIMIN 24-7 COPPER are long acting boluses for the oral administration of copper to cattle. The boluses dissolve in the rumen, releasing the copper oxide needles. AGRIMIN 24-7 COPPER are designed to maintain copper levels over the whole grazing season (approx. 6 months). Each bolus contains 20 g of copper oxide needles, to supply 16.8 g elemental copper.

AGRIMIN 24-7 COPPER are indicated for the prevention and treatment of copper deficiency in cattle over 200kg liveweight.

**WITHHOLDING PERIODS:**  
MEAT: Zero (0) days  
MILK: Zero (0) days

**TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI):**  
ESI from Pacific Biologics on (07) 3283 5077 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues))

**SAFETY DIRECTIONS**  
May irritate the eyes and skin. Avoid contact with the eyes and skin. Wash hands after use.

**FIRST AID**  
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

**DISPOSAL**  
Dispose of container by wrapping with paper and putting in garbage.

**STORAGE**  
Store below 25°C (air conditioning). Store in original container.

Protect from light.

Dose animals early winter unless copper supplementation is required immediately.

In areas where liver fluke are endemic, treat animals with a flukicide at least 3 weeks before using AGRIMIN 24-7 COPPER boluses.

The boluses may change shape if exposed to high temperatures (above 40°C).

**WITHHOLDING PERIODS:**

**MEAT: Zero (0) days**

**MILK: Zero (0) days**

**TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI):**

ESI from Pacific Biologics on (07) 3283 5077 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues))

**SAFETY DIRECTIONS**

May irritate the eyes and skin. Avoid contact with the eyes and skin. Wash hands after use.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

**DISPOSAL**

This product is contraindicated for use in cattle suffering from jaundice and other liver disorders such as liver fluke, or stock grazing plants containing toxic alkaloids, e.g. Heliotrope, Ragwort, Patersons Curse (Salvation Jane).

Seek the advice of a veterinary or cattle husbandry specialist to ascertain copper deficiency and type of treatment required for cattle under consideration.

**Dosage and Administration**

Only repeat treatment within 12 months when specifically indicated by laboratory tests. Administer orally. Place the bolus at the back of the mouth using a suitable applicator. It is important that the applicator's instructions are strictly followed. One bolus should be administered per animal over 200 kg. One bolus per 200 kg.

**General Directions**

To be effective, the bolus must be swallowed by the animal.  
Dose pregnant females early in pregnancy to allow protection of the calf against copper deficiency.

**Contraindications**

SHOULD NOT BE ADMINISTERED TO CATTLE LESS THAN 5 MONTHS OF AGE OR 200 kg BODYWEIGHT

Excessive copper is toxic. This product is contraindicated for use unless copper deficiency is confirmed.

**DIRECTIONS FOR USE**

AGRIMIN 24-7 COPPER are long acting boluses for the oral administration of copper to cattle. The boluses dissolve in the rumen, releasing the copper oxide needles. AGRIMIN 24-7 COPPER are designed to maintain copper levels over the whole grazing season (approx. 6 months). Each bolus contains 20 g of copper oxide needles, to supply 16.8 g elemental copper.

AGRIMIN 24-7 COPPER are indicated for the prevention and treatment of copper deficiency in cattle over 200kg liveweight.

24 x 23 g Boluses

SUSTAINED RELEASE BOLUSES FOR CATTLE

Copper 20g



SUSTAINED RELEASE BOLUSES FOR CATTLE

24 x 23 g Boluses

Copper 20g



READ SAFETY DIRECTIONS BEFORE OPENING OR USING FOR ANIMAL TREATMENT ONLY



Copper 20g

SUSTAINED RELEASE BOLUSES FOR CATTLE

ACTIVE CONSTITUENT: 732 g/kg Copper (as copper oxide)

Each bolus contains 20 g copper oxide needles, to supply 16.8 g elemental copper. For the prevention and treatment of copper deficiency in cattle over 200 kg bodyweight.

24 x 23 g Boluses

cattle



Copper 20g

SUSTAINED RELEASE BOLUSES FOR CATTLE

24 x 23 g Boluses